

Result Navigator

Positive Test Result: *EPCAM*

Positive test results identify a change, or “misspelling,” of DNA that is known or predicted to cause an increased risk for cancer. DNA is the blueprint of life—it carries all of the genetic information needed for our bodies to function. A section of DNA with a specific job is called a gene. A pathogenic variant is a change in a gene that is known to cause increased risk for cancer. A likely pathogenic variant is a change in a gene that is very likely to cause increased risk for cancer. Another name for these types of variants is mutation.

After a positive test result, there can be many questions about what to do next.

This guide is a supplement to Riscover™ Hereditary Cancer test results, helping patients and healthcare providers understand the results and together determine the best path forward.

Navigate Your Results



Learn | page 2

Review the test results along with this guide, and discuss the meaning of these results with your healthcare provider.



Plan | page 3

Develop a healthcare plan for early detection and prevention of specific cancers.



Share | page 4

Understand the risks for other family members, and consider how to inform relatives of their risks.



Connect | page 5

Reach out to medical experts and find local and national groups that can provide more information and support.

Glossary of Terms

CA-125: A substance that can indicate the development of ovarian cancer. Testing for CA-125 is performed on a sample of blood.

Colonoscopy: Visual exam of the inside of the colon with a flexible, lighted tube inserted through the rectum.

EGD (Esophagogastroduodenoscopy): Also called upper endoscopy, this is an endoscopic test that examines the esophagus (part of the gastrointestinal tract that connects the throat to the stomach), stomach, and first part of the small intestine (called the duodenum).

Gene: A section of DNA with a specific job.

H. pylori: A type of bacteria found in the stomach that can cause ulcers and also increases the risk of stomach cancer.

Hepatobiliary tract: Includes the liver, gall bladder, and bile ducts.

Hysterectomy: Surgical removal of the uterus.

Mutation: A change in a gene that causes or is capable of causing disease. Another word for pathogenic/likely pathogenic variants.

Pathogenic: Causing or capable of causing disease.

Risk: The chance, or possibility, of developing a disease.

Salpingo-oophorectomy: Surgical removal of the ovaries and fallopian tubes.

Transvaginal ultrasound: An imaging test performed by inserting an ultrasound probe into the vagina to take detailed images of the uterus, ovaries, and fallopian tubes.

Urinalysis: A test that measures different substances in the urine that may indicate a wide range of disorders such as kidney disease or the development of cancer.

Urinary tract: The system in the body that eliminates waste through urine. The urinary tract includes the kidneys, ureters (part of the body that takes the waste from the kidneys and transports it to the bladder), bladder, and urethra (part of the body that moves urine outside of the body).

Variant: A change, or “misspelling,” in the DNA sequence. Variants can be benign (not associated with disease), pathogenic (associated with disease), or of uncertain significance.



Your Diagnosis: Lynch Syndrome

Lynch syndrome is an inherited disorder caused by a pathogenic variant in one of the following mismatch repair genes: *MLH1*, *MSH2*, *MSH6*, or *PMS2*. Certain pathogenic variants in a fifth gene, *EPCAM*, can also cause Lynch syndrome by “turning off” *MSH2*. The normal job of a mismatch repair gene is to check newly formed DNA for errors. When these genes have pathogenic variants, new errors in DNA can go uncorrected, which can lead to the development of cancer. Lynch syndrome is associated with an increased lifetime risk of colorectal cancer. Females with Lynch syndrome are also at risk for endometrial (uterine) and ovarian cancers. Males and females are also at an increased risk for cancers of the stomach, hepatobiliary tract, urinary tract, small bowel, and brain/central nervous system.

Lynch syndrome accounts for approximately 3 – 5% of all colorectal cancers. About 1 in 400 people in the general population have Lynch syndrome. People who have two pathogenic variants (one inherited from each parent) in the same gene are affected with a condition called constitutional mismatch repair deficiency (CMMRD). The risks described below are for people with one pathogenic variant.


Your Risk

Not all people with positive test results will develop cancer, but the chance for cancer is higher than the general population risk. The graphs below show how much the risk for cancer increases when a pathogenic variant is found. The cancer risks are presented in ranges based on multiple studies of families who have a pathogenic variant. There is not enough evidence available to give a specific risk to a person or family.

LIFETIME CANCER RISKS

■ General population

■ People with *EPCAM* positive results



Your Screening and Management Options

Listed below are the intervention options that are recommended by medical experts for individuals who carry this cancer gene variant. Your healthcare provider can explain these options in more detail, and together you can make a plan for cancer prevention and/or early detection.

COLORECTAL CANCER

Colonoscopy	A colonoscopy every 1 – 2 years is recommended starting at age 20 – 25 (or 2 – 5 years before the earliest diagnosis in the family if under age 25).
Aspirin	Your healthcare provider may prescribe aspirin to help prevent colorectal cancer, based on your individual situation.

ENDOMETRIAL/OVARIAN CANCERS (FEMALES)

Consider total hysterectomy and bilateral salpingo-oophorectomy	Removal of the uterus, ovaries and fallopian tubes, should be considered after a woman has had her last child.
Consider transvaginal ultrasound, endometrial sampling, and CA-125 (at clinician discretion)	Your healthcare provider may consider an annual transvaginal ultrasound to screen for endometrial and ovarian cancers. They may also consider endometrial biopsy every 1 – 2 years and/or a blood test for cancer antigen (CA-125).
Dysfunctional uterine bleeding warrants evaluation	Seek evaluation with your healthcare provider when present.

STOMACH/SMALL BOWEL CANCERS

EGD with extended duodenoscopy	Consider an EGD with extended duodenoscopy every 3 – 5 years beginning at age 30 – 35, based on your personal or family history of stomach cancer and your ethnicity. (Stomach cancer in Lynch syndrome is more common in people of Asian descent.)
Consider testing for and treating <i>H. pylori</i>	Consider testing for <i>H. pylori</i> as recommended by your healthcare provider.

HEPATOBIILIARY TRACT CANCER

No specific guidelines exist	Your healthcare provider can recommend screening options based on your personal and family history of hepatobiliary tract cancer.
------------------------------	---

URINARY TRACT CANCER

Urinalysis	Consider an annual urinalysis beginning at age 30 – 35.
------------	---

BRAIN/CNS CANCER

Physical and neurological exam	Consider an annual physical and neurological exam beginning at age 25 – 30.
--------------------------------	---

PANCREATIC CANCER

No specific guidelines	Your healthcare provider can recommend screening options based on your personal and family history of pancreatic cancer.
------------------------	--


Your Family Matters

Cancer gene variants are inherited in an autosomal dominant manner. This means that when a parent tests positive, each child has a 1 in 2 chance to inherit the variant. Some people who inherit the variant will develop cancer, and some will not, but the risks are significantly increased compared to the average person. People in the same family may develop different types of cancers at varying ages, and the pattern of cancer and age of onset can differ from family to family.

Gene variants can be passed through families for many generations. When a gene variant is identified in one person in the family, relatives can be tested for just that variant. Sharing information about test results with family members is important, because it allows them to make informed choices about their healthcare and to decide whether they want genetic testing.

Some people who have pathogenic and likely pathogenic variants are interested in learning how to avoid passing these variants on to their children. If you are planning to have a child, you may wish to consult with a reproductive medicine specialist to discuss all of your reproductive options, including use of an egg or sperm donor, preimplantation genetic diagnosis, and prenatal genetic testing.

The likelihood for family members to carry the same gene variant depends on how closely related they are to the person with a known gene variant.



Your Expert Team

As you take your next steps, it can help to have a team of experts on your side. This team may include your primary care provider and a number of specialists who will help along the way.

If you haven't already done so, you may wish to speak with a genetic counselor. Genetic counselors are healthcare professionals with specialized training in medical genetics and counseling. They can help you understand your test results, discuss your healthcare plan, and connect to other resources in your area. Your healthcare provider may be able to provide a referral to a local genetic counselor. You can also find a genetic counselor through the National Society of Genetic Counselors at www.nsgc.org.

For questions about your test results, Progenity's genetic counseling team is available to help. To reach a genetic counselor, call +1 855-293-2639 and select option 3.

Helpful Resources

GeneReviews

ncbi.nlm.nih.gov/books/NBK1211

A detailed clinical summary of Lynch syndrome written for healthcare providers.

Hereditary Colon Cancer Takes Guts

hctakesguts.org/about

A support group whose mission is to serve the hereditary colon cancer community by connecting individuals to resources and promoting research and health care initiatives.

Lynch Syndrome International

lynchcancers.com

A support group whose mission is to provide support to those with Lynch syndrome and to promote public awareness, education, and research.

Lynch Syndrome Screening Network

lynchscreening.net

A support group for medical providers whose mission is to promote universal Lynch syndrome screening on colorectal and endometrial cancers and provide resources for appropriate management guidelines.

National Comprehensive Cancer Network (NCCN)

nccn.org

Expert guidelines for cancer screening and treatment. Versions available for both providers and patients.

REFERENCES

Kohlmann W and Gruber SB. *Lynch Syndrome*. 2004 Feb 5 [Updated 2014 May 22]. In: Pagon RA, Adam MP, Ardinger HH, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2016. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK1211>

National Comprehensive Cancer Network. Genetic/Familial High-Risk Assessment: Colorectal (Version 1.2017). Available from: <http://www.nccn.org>

5230 S. State Road, Ann Arbor, MI 48108 USA • Tel +1 855-293-2639 • progenity.com

Progenity is a CLIA-certified clinical laboratory and is accredited by the College of American Pathologists (CAP). The information contained in this document is provided by Progenity as an educational service for clinicians and their patients. © 2015, 2016 Progenity, Inc. All rights reserved. Progenity® is a registered service mark of Progenity, Inc. Riscover™ is a trademark of Progenity, Inc. WH-55006-01 REV 072017